

Dosage Chart Tylenol / Ibuprofen

Acetaminophen (dose for Children with fever) (e.g., Tylenol, Store Brand Acetaminophen) dosage based on weight.						Ibuprofen dosages (use Children's oral liquid 100mg/tsp (e.g. Motrin, Advil, Store Brand Ibuprofen) do not use for babies less than 6 months.			
Childre n Age	Appro x Weight (Lbs)	Infant Oral Suspension 160mg/5(ml)) dose-in (ml) **	Childrens' Liquid-dose 160mg/5(ml)) dose in (Tsp)	Children's Chewable 80mg Tablets	Junior Chew Tablet s 160 mg	Childre n Age	Appro x Weight (Lbs)	Dose (mg) 100m g /tsp	Dose (teas)
Under 4 Months	8 12	1.25 2.5ml	1/4 1/2	—	—	6 Months 11 Months	13 17	50	½
4 Months 11 Months	12 18	2.5ml 3.75	1/2 3/4	1 1 ½	—	12 Months 23 Months	18 23	100	1
12 Months 23 Months	18 24	3.75 5	3/4 1	1 ½ 2		2 Years 3 Years	24 35	150	1 ½
2 Years 3 Years	24 36	5	1 1 ½	2 3	1 1 ½	4 Years 5 Years	36 47	200	2
4 Years 5 Years	36 47	— —	1 ½ 2	3 4	1 ½ 2	6 Years 8 Years	48 59	250	2 ½
6 Years 8 Years	48 60	— —	2 2	4 4	2 2	9 Years 10 Years	60 71	300	3
9 Years 10 Years	60 72	— —	2 ½ 3	5 6	2 ½ 3	11 Years 12 Years	72 95	400	4
11 Years	72 92	— —	3 4	6 8	3 4				
12 Years	96 & over	— —	4	8	4				
Acetaminophen -One dose may be given every 4 to 6 hrs as needed. ** Use syringe provided						Ibuprofen -One dose may be given every 6 to 8 hrs as needed. Do not use with active vomiting, a head injury or where there is active bleeding.			